

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search	PubMed	for				Go	Clear
		Limits	Preview/Index	History	Clipboard		

Display	Abstract	Save	Text	Order	Add to Clipboard
---------	----------	------	------	-------	------------------

Entrez
PubMed☐ 1: *J Endocrinol* 1999 Sep;162(3):371-9

Related Articles, Books, LinkOut

[JOE Online](#)PubMed
Services**Effects of dexamethasone on proliferation, activity, and cytokine secretion of normal human bone marrow stromal cells: possible mechanisms of glucocorticoid-induced bone loss.****Kim CH, Cheng SL, Kim GS**

Division of Endocrinology, Department of Internal Medicine, Soonchunhyang University College of Medicine, Seoul, Korea.

NEW

Related
Resources

It is well documented that glucocorticoid excess causes bone loss, but the mechanisms of these effects remain poorly defined. To understand further the mechanisms of glucocorticoid-induced osteoporosis, we investigated the effects of glucocorticoids on bone formation and bone resorption by examining the proliferation, functional activities, and cytokine secretion of cultured human bone marrow stromal cells (hBMSC). Treatment with dexamethasone for 24 h at the concentration of 10^{-8} M significantly suppressed $[(3)H]$ thymidine incorporation and further inhibition was observed with longer treatment (8 days) or higher concentration (10^{-7} M). Alkaline phosphatase activity of hBMSC was markedly stimulated with addition of dexamethasone (10^{-8} M), to $191 \pm 22\%$ (after 4 days) and $317 \pm 46\%$ (after 7 days) of control. Dexamethasone (10^{-8} M) treatment for 48 h decreased the incorporation of $[(3)H]$ proline into collagenase-digestible protein (CDP; $43.7 \pm 7.9\%$ of control) and non-collagen protein ($65.2 \pm 8.4\%$ of control), with a greater effect on CDP. Northern blot analysis indicated that $\alpha 1(I)$ -collagen mRNA level was decreased by dexamethasone to $27.6 \pm 9.0\%$ of the control value after 1 day of exposure, and to $55.2 \pm 6.2\%$ after 7 days. Dexamethasone markedly suppressed basal production of interleukin (IL)-6 and IL-11 and that stimulated by parathyroid hormone (PTH), IL-1 α , or tumour necrosis factor- α in a dose-dependent manner. These results suggest that the glucocorticoid-induced bone loss is derived at least in part via inhibition of bone formation, which includes the suppression of osteoblast proliferation and collagen synthesis. As both basal and PTH-stimulated production of IL-6 and IL-11 are decreased by dexamethasone, the increased bone resorption observed in glucocorticoid-induced osteopenia does not appear to be mediated by IL-6 or IL-11.

PMID: 10467228, UI: 99398450